

Cluster deprojection with joint lensing, X-ray, and Sunyaev-Zeldovich data

Katrin Reblinsky and Matthias Bartelmann

Max Planck Institut für Astrophysik, P.O. 1523, D-85740 Garching, Germany

Abstract. We propose a new cluster deprojection algorithm for recovering cluster structure along the line of sight (los) based on the Richardson-Lucy (RL) algorithm, which recovers a nonnegative theoretical distribution from a given projection. To optimize our reconstruction we combine X-ray and Sunyaev-Zeldovich (SZ) data with gravitational lensing maps. The combination of these three data types constrains the structure of rich clusters of galaxies. To validate our new approach, we apply the algorithm to gas-dynamical cluster simulations. A reconstruction using the lensing potential is shown.

1. Richardson-Lucy deprojection algorithm

Resolving the los-structure of 3d objects can be formulated as inversion problem,

$$\psi(x) = \int \phi(\zeta) P(x|\zeta) d\zeta,$$

where $\phi(\zeta)$ is the function of interest, $\psi(x)$ is the function accessible through measurement, and the integral kernel $P(x|\zeta)$ is normalized and non-negative as $\psi(x)$ and $\phi(\zeta)$ represent probability distribution functions.

Using Bayes theorem for conditional probabilities, a two step *iterative reconstruction algorithm* (Lucy, 1974; Lucy, 1994)

$$\psi^r(x) = \int \phi^r(\zeta) P(x|\zeta) d\zeta \quad \phi^{r+1}(\zeta) = \phi^r(\zeta) \int \frac{\psi(x)}{\psi^r(x)} P(x|\zeta) dx$$

for approximating $\phi(\zeta)$ can be derived, with ϕ^{r+1} being the $(r+1)$ -st estimate. For systems with axial symmetry inclined with respect to the los an explicit expression for $P(x|\zeta)$ was derived by Binney et al. (1990).

2. The proposed algorithm

The three observables depend differently on the los-structure of the 3d gravitational potential (Reblinsky & Bartelmann (1999)):

$$\text{Lensing potential} \quad \psi(x, y) \propto \int_{-\infty}^{+\infty} \phi(R, Z) dz,$$




Figure 1. Reconstruction of the 3d gravitational potential ϕ (displayed is $-\phi$) from the lensing potential. Inclination angle $i = 30^\circ$. Left: the result after 7 iteration cycles; Right: the original cluster potential used to compute the input data, i.e. the lensing potential ψ .

$$\text{X-ray emission} \quad S_x(x, y) \propto \int_{-\infty}^{+\infty} \sqrt{T} \exp[-2\phi'(R, Z)] dz,$$

$$\text{SZ decrement} \quad \Delta T_{\text{SZ}}(x, y) \propto \int_{-\infty}^{+\infty} T \exp[-\phi'(R, Z)] dz.$$

For the X-ray case and the SZ-effect an isothermal gas distribution is assumed. The inclination angle i and the temperature T are input parameters. This leads to the following *combined two step reconstruction formula*:

$$\text{Step 1:} \quad F_{n_i} = \int f_{n_i}(x, y) P(x, y | R, Z) dx dy$$

$$\text{Step 2:} \quad \phi_{r+1} \propto \phi_r \left[\alpha F_{n_1} + \beta \left(1 - \frac{1}{2} \ln F_{n_2} \right) + \gamma \left(1 - \ln F_{n_3} \right) \right],$$

where the f_{n_i} are defined as $f_{n_1} = \frac{\psi}{\psi_n}$, $f_{n_2} = \frac{S_x}{S_{x,n}}$, and $f_{n_3} = \frac{\Delta T_{\text{SZ}}}{\Delta T_{\text{SZ},n}}$.

3. Application to test data from gas-dynamical cluster simulations

Using the lensing potential $\psi(x, y)$ as input data, the characteristic features of the true 3d potential $\phi(R, Z)$ are recovered very well after 7-8 iteration cycles. We also find that the reconstruction is insensitive to the initial guess, even $\phi(R, Z) = \text{const}$ can be used as in Fig. 1.

Acknowledgments. This work was supported in part by the Sonderforschungsbereich 375 of the Deutsche Forschungsgemeinschaft.

References

- Binney J.J., Davies R.L., Illingworth G.D., 1990, ApJ, 361, 78
- Lucy L.B., 1974, AJ, 79, 745
- Lucy L.B., 1994, A&A, 289, 983
- Reblinsky K., Bartelmann M., 1999, in preparation